

The Graphic Reality of Artificial Immunization



Raymond Obomsawin, Ph.D.
Senior Advisor on First Nations Health
National Aboriginal Health Organization
Ottawa, Canada


November, 2010

Independent Updates – March, 2015 & Sept, 2019


Natural Infectious Disease Declines Preceding Mass Public Artificial Immunization Efforts

Graphs one (1) through ten (10) reveal that in North America, Europe, and the South Pacific, major declines in life-threatening infectious diseases occurred historically either without, or many decades in advance of mass immunization efforts for specific diseases as listed. This represents irrefutable evidence that improvements in nutrition, hygiene, sanitation, & safe drinking water were the actual determinants underlying the historically massive decline of a wide range of infectious diseases throughout Western World nations.

**United States
Tuberculosis Mortality
Rates per 100,000 Infants (1900-1960)**




Source: John H. Dingle; Life and Death in Medicine; Scientific American; 1973; p. 56.




Source: Data derived from - Vital Statistics of the United States 1937-1960; and Historical Statistics of the United States: Colonial Times to 1970 Part 1 Ch. B Vital Statistics and Health and Medical Care, pp. 44-86H.

United States Annual Influenza Mortality Rates per 100,000 (1933-1965)




Source: Doshi, P., Trends in Recorded Influenza Mortality: United States 1900-2004, American Journal of Public Health, May 2008, vol. 98, no. 5, p. 941.

United States Mean Annual Pertussis Mortality Rates Per 100,000 (1918-1960)




Source: Data derived from: Vital Statistics of the United States 1937-1960; and Historical Statistics of the United States: Colonial Times to 1970 Part 1 Ch. B Vital Statistics and Health and Medical Care, pp. 44-86H.

5. England & Wales Mean Annual Pertussis Mortality Cases Children under 15 (1850-1965)



Source: Thomas McKeown, *The Role of Medicine: Dream, Mirage or Nemesis?*; Basil Blackwell; Oxford, UK; 1979; p. 103

**England & Wales
Mean Annual Measles Mortality
Cases Children under 15 (1850-1965)**




**Measles
Vaccination Begins in
England & Wales**

Source: McKeown, T., *The Role of Medicine: Dream, Mirage or Nemesis?*; Basil Blackwell; Oxford, UK; 1979; p. 105; & Waltzkin, H., in *The Relevance of Social Science for Medicine*; Springer; 1st edition, Dec. 31, 1980


England

Scurvy (Severe Vitamin C Deficiency) & Measles - *Parallel Decline in Mortality Rates per 100,000 (1919-1967)*




England

Scurvy (Severe Vitamin C Deficiency) & Pertussis - *Parallel Decline in Mortality Rates per 100,000 (1919-1967)*




Canada Tuberculosis Mortality Rates per 100,000 (1880-1960)



Source: based on data at: Timeline of TB in Canada <http://www.lung.ca/tb/tbhistory/timeline/>; <http://www.thecanadianencyclopedia.com/index.cfm?PgNm=TCE&Params=A1ARTA0008151>
Public Health Agency of Canada: <http://www.phac-aspc.gc.ca/publicat/cig-gci/p04-bcg-eng.php>; &
PHAC on BCG usage in Canada: http://www.phac-aspc.gc.ca/tbpc-latb/bcgvac_1206-eng.php

New Zealand Tuberculosis Mortality Rates Per Million (1880-1960)




Source: Director General Annual Mortality Reports Covering 1872-1960, New Zealand Parliamentary Journals for the Years Specified.

Actual Influenza Incidence when Flu-like Respiratory Infections Occur

Graph eleven (11) shows that when Flu-like illnesses occur, 90% of the time it is typically not the Flu. This means that assuming widely employed Flu vaccines were actually effective, 90% of recipients would remain unprotected from the multiple viruses associated with a diverse range of Flu-like respiratory infections.

Flu-Like Acute Respiratory Infections




Nicholson, K.G. et. al., Acute viral infections of upper respiratory tract in elderly people living in the community: comparative, prospective, population based study of disease burden, British Medical Journal, No. 315, October 25, 1997, pp. 1060-1064

Artificial Immunization Efficacy

Graphs twelve (12) through twenty (20) clearly show that the practice of artificial immunization is not by any means a proven and foolproof measure for protection from various infectious disease conditions. It is often inconsequential epidemiologically, and in some cases it demonstrably worsens infectious disease health outcomes.

Children Under 2 Yrs of Age
Inactivated Influenza Vaccine



A 3D pie chart with a single white slice representing 0% of the total. The chart is set against a blue gradient background. The text '0% Effective' is centered on the white slice.

0%
Effective

**Source: Cochrane Collaboration Database of Systematic Reviews, (John Wiley & Sons, Ltd.)
2006 (1) Article No. CD004879 – Covers 51 Studies on 260,000 children**

Elderly Living in Communities
& Group Homes
Inactivated Influenza Vaccine



**Negligible or 0%
Effectiveness**

Source: Cochrane Collaboration Database of Systematic Reviews, (John Wiley & Sons, Ltd.) 2006 (3)
Article No. CD004876 – Covers 64 Studies, over 40 years of influenza vaccination and see:
<http://www.bmj.com/cgi/content/full/333/7574/912>

BCG for Tuberculosis

Note: Tuberculosis found higher
among two (2) dose Vaccinated
versus Placebo Group

0%
Effective

Source: Randomised controlled trial of single BCG, repeated BCG, or combined BCG and *killed Mycobacterium leprae* vaccine for prevention of leprosy and tuberculosis in Malawi; The Lancet, Volume 348, Issue 9019, Pages 17 - 24, July 6, 1996.

BCG for Tuberculosis


Note: In years 0-2.5 the vaccinated
had double the incidence of
Tuberculosis versus Placebo Group

0%
Effective

Source: Double blind randomized controlled trial of BCG's effectiveness on 250,000 subjects Tuberculosis Research Centre (ICMR), Chennai, India: Indian Journal of Medical Research, 110, August 1999, pp. 56-69.

Mumps Outbreak in Highly Vaccinated Population


2006
Iowa



Source: Center for Disease Control ,
MMWR 55 (20); May 26, 2006; pp. 559-63.

Chickenpox Outbreak in Highly Vaccinated Population


2001
Oregon



Source: Pediatrics - Vol. 113; No. 3;
pp. 455-459; (2004)

Pertussis Outbreak in Highly Vaccinated Population


1993
Ohio



Source: N.Z. Miller; *Vaccine Safety Manual*;
N.A. Press, Sante Fe, New Mexico; p. 140; (2008)
(Refers to CDC & Official Surveillance data)

Measles Outbreak in Highly Vaccinated Population


1985
Texas



Source: New England Journal of Medicine -
Vol. 316; No. 13; pp. 771-774; (1987)

Measles Outbreak in Highly Vaccinated Population

1988
Colorado




Source: “A Measles Outbreak at a College with a Pre-matriculation Immunization Requirement” - American Journal of Public Health, March 1991, Vol. 81, No. 3

Artificial Immunization Documented Dangers

Graphs twenty one (21) through thirty (30) reveal that increases in the number of mandated vaccinations correlate with a major rise in infant mortality (i.e. death rates) among children under age 5, and infants.


Artificial immunization is also linked to Sudden Infant Death Syndrome; various degenerative diseases, including Diabetes; and appears to cause general immune system impairment. Evidence also points to the practice as a principal factor in the recent massive increases in neurodegenerative conditions, including unprecedented Autism levels in children.

COUNTRIES & NUMBER OF VACCINES MANDATED UP TO AGE 5 *UNDER AGE FIVE MORTALITY RATES - 2007*




Under Age 5 Mortality statistics: http://www.who.int/whosis/whostat/EN_WHS09_Table1.pdf
WHO – World Health Statistics 2009 & Govt. Vaccines figures: Generation Rescue Inc. 2009
<http://www.generationrescue.org/documents/SPECIAL%20REPORT%20AUTISM%202.pdf>

COUNTRIES & NUMBER OF VACCINES MANDATED UP TO AGE ONE *INFANT MORTALITY RATES - 2009*




UNDER AGE 5 INFLUENZA DEATHS BEFORE & AFTER U.S. CDC MANDATES FLU VACCINES IN EARLY CHILDHOOD




Under Age 5 Influenza Mortality statistics derived from: Center for Disease Control Vital Statistics Reports covering Years 1999-2003 reported in Miller, N.Z., Vaccine Safety Manual, New Atlantean Press, Sante Fe, New Mexico, 2008, p. 97.

PERTUSSIS VACCINE & *SUDDEN INFANT DEATH SYNDROME*



2/3 of 103 infants had been vaccinated with pertussis prior to death, of which 6.5% died within 12 hours; 13% within 24 hours; 26% within 3 days; 37%, 61% & 70% within 1, 2, & 3 weeks respectively. Source: Torch W., Neurology - 32 (4 – Pt. 2) A, 1982, pp. 169-170.


Measles Vaccination & Inflammatory Bowel Diseases



Source:The Lancet - Vol. 345; 8957; 1995, pp. 1062-1063.


Average Incidence First Five (5) years of Life

Nederlands Vereniging Kritisch Prikken 2004 Survey Findings




Absolute Incidence N=543

Nederlands Vereniging Kritisch Prikken 2004 Survey Findings




Absolute Incidence (Non-Vaccinated in Relation to Vaccinated to N = 312 Per Group)


BCG Vaccination Mandated in Schools & Diabetes Rates by Country



Cumulative Incidence IDDM/1,000,000 UK



Source: Journal of Pediatric Endocrinology & Metabolism, 16, pp. 495-508; (2003)



Source: Adapted from Blaxil, M., Vaccine Mercury Burden & Autism Risk (US) IOM 7/2001,
http://www.healing-arts.org/children/marksslide_files/slide0003.htm